

FAZE RAZVOJA ULJANE REPICE

HERBICIDI	KOROVI	prije sjetve	sjetva	prije nicanja	kotiledoni	2 prava lista	rozeta	porast lisne mase	oblikov. pupova	puno cvjetanje	biološka zrioba
 GLIFOL	svi iznikli korovi prije obrade	3- 4 l/ha									
 PIKOGAL PLUS	jednogodišnji i višegodišnji širokolisni i neki uskolisni korovi					0.2 – 0.3 l/ha					
 RAFAL 120* *Ima registraciju u EU.	iznikli jednogodišnji uskolisni korovi uključujući i divlji sirak iz rizoma i pirevinu						1 – 2 l/ha				
FUNGICIDI	BOLESTI										
 AKORD* *Ima registraciju u EU.	suva trulež korijena, koncentrična pjegavost, bijela i siva trulež						0.1 l/ha na svaki list uljane repice (6 listova = 0.6 l/ha)				
 PROPULSE 250 SE	siva, bijela i suva trulež, crna pjegavost, pepelnica						1 l/ha				
INSEKTICIDI	ŠTETOČINE										
 POLUX ili DECIS EC 100	repičina lisna osa, repičin buvač, repičin sjajnik, repičina pipa, podgrizajuće sovice					0,3 l/ha ili 0,075 l/ha					