

FAZE RAZVOJA STRNIH ŽITA

HERBICIDI	KOROVI	prije sjetve	prije nicanja	faza 2.-3. lista	početak bokorenja	početak vlatanja	faza 1. koljenca	faza 2. koljenca	klasanje	cvjetanje
 SEKATOR OD	iznikli jednogodišnji i višegodišnji širokolisni korovi uključujući i bročiku (Galium)						0,1 – 0.15 l/ha			
 HUSSAR OD	iznikli jednogodišnji i višegodišnji uskolisni i širokolisni korovi						0,1 l/ha			
 XANADU	iznikli jednogodišnji i višegodišnji širokolisni korovi uključujući i bročiku (Galium)						0,1 kg/ha			
 BONACA EC	iznikli jednogodišnji i višegodišnji širokolisni korovi uključujući i bročiku (Galium)						0,4 – 1,2 l/ha			
 TEZIS	iznikli jednogodišnji i višegodišnji širokolisni korovi						10 g/ha			
FUNGICIDI	BOLESTI									
 RANCONA 15 ME	glavnica pšenice, fuzariozna palež kljanaca i trulež korjena strnih žita, sniježna plijesan žita, otkrivena gar ječma	100 - 133 ml na 100 kg sjemena								
 AKORD	pepelnica, rđa, pjegavost lista						1 l/ha			
 FALCON FORTE 460 EC	pepelnica, rđa, siva pjegavost lista i klasa pšenice, mrežasta pjegavost ječma							0,7 – 1 l/ha		
 PROSARO 250 EC	pepelnica, rđa, pjegavost lista i klasa, fuzarioze klasa								0.85 – 1 l/ha	
 ANTRE PLUS	pepelnica, rđa, mrežasata pjegavost ječma, pjegavost lista i pljevica								1, 5 l/ha	
 NEOTOPSIN 70 WG	rđa, zemljišni patogeni									0,9 kg/ha
INSEKTICIDI	ŠTETOČINE									
 POLUX ili DECIS 100 EC* *Ima registraciju u EU.	žitna pijavica (lema), lisne vaši						0,2 - 0,3 l/ha ili 0,075 – 0,125 l/ha			
 KOZAK* *Ima registraciju u EU.	žitna pijavica (lema), lisne vaši						0,15 l/ha			